Journal of Knot Theory and Its Ramifications
© World Scientific Publishing Company

arXiv:1512.05481v2 [math.GT] 3 Mar 2016

The volume of hyperbolic cone-manifolds of the knot with Conway's notation $C(2n, 3)$

Ji-Young Ham *
Department of Science, Hongik University, 94 Wausan-ro, Mapo-gu, Seoul, 04066, Department of Mathematical Sciences, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Korea
jiyoungham1@gmail.com

Joongul Lee †
Department of Mathematics Education, Hongik University, 94 Wausan-ro, Mapo-gu, Seoul, 04066, Korea
jglee@hongik.ac.kr

ABSTRACT

Let $C(2n, 3)$ be the family of two bridge knots of slope $(4n+1)/(6n+1)$. We calculate the volumes of the $C(2n, 3)$ cone-manifolds using the Schläfli formula. We present the concrete and explicit formula of them. We apply the general instructions of Hilden, Lozano, and Montesinos-Amilibia and extend the Ham, Mednykh, and Petrov's methods. As an application, we give the volumes of the cyclic coverings over those knots. For the fundamental group of $C(2n, 3)$, we take and tailor Hoste and Shanahan's. As a byproduct, we give an affirmative answer for their question whether their presentation is actually derived from Schubert's canonical 2-bridge diagram or not.

Keywords: hyperbolic orbifold, hyperbolic cone-manifold, volume, $C(2n, 3)$, orbifold covering, Riley-Mednykh polynomial.

Mathematics Subject Classification 2010: 57M25, 57M27

1. Introduction

Let us denote $C(2n, 3)$ by T_{2n} and the exterior of T_{2n} by X_{2n}. By Mostow-Prasad rigidity, X_{2n} has a unique hyperbolic structure. Let ρ_∞ be the holonomy representation from $\pi_1(X_{2n})$ to $PSL(2, C)$ and denote $\rho_\infty(\pi_1(X_{2n}))$ by Γ, a Kleinian group. X_{2n} is a $(PSL(2, C), \mathbb{H}^3)$-manifold and can be identified with $\mathbb{H}^3/\Gamma$. Thurston's orbifold theorem guarantees an orbifold, $X_{2n}(\alpha)$, with T_{2n} as the singular locus

*This work was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (No. NRF-2008-341-C00004 and NRF-R01-2008-000-10052-0).
†This work was supported by 2015 Hongik University Research Fund.

and the cone-angle $\alpha = 2\pi/k$ for some nonzero integer k can be identified with $\mathbb{H}^3/\Gamma'$ for some $\Gamma' \in PSL(2,C)$; the hyperbolic structure of X_{2n} is deformed to the hyperbolic structure of $X_{2n}(\alpha)$. For the intermediate angles whose multiples are not 2π and not bigger than π, Kojima [11] showed that the hyperbolic structure of $X_{2n}(\alpha)$ can be obtained uniquely by deforming nearby orbifold structures. Note that there exists an angle $\alpha_0 \in [\frac{2\pi}{3}, \pi)$ for each T_{2n} such that $X_{2n}(\alpha)$ is hyperbolic for $\alpha \in (0, \alpha_0)$, Euclidean for $\alpha = \alpha_0$, and spherical for $\alpha \in (\alpha_0, \pi]$ [18,8,11,19]. For further knowledge of cone manifolds you can consult [1,7].

Explicit volume formulae for hyperbolic cone-manifolds of knots and links are known a little. The volume formulae for hyperbolic cone-manifolds of the knot 4_1 [8,11,12,15], the knot 5_2 [13], the link 5_1^2 [16], the link 6_2^2 [17], and the link 6_3^2 [2] have been computed. In [9] a method of calculating the volumes of two-bridge knot cone-manifolds were introduced but without explicit formulae. In [7], explicit volume formulae for hyperbolic twist knot cone-manifolds are computed. Similar methods are used for computing Chern-Simons invariants of twist knot orbifolds in [6].

The main purpose of the paper is to find an explicit and efficient volume formula for hyperbolic cone-manifolds of the knot T_{2n}. The following theorem gives the volume formula for $X_{2n}(\alpha)$. Note that if $2n$ of T_{2n} is replaced by an odd integer, then T_{2n} becomes a link with two components. Also, note that the volume of hyperbolic cone-manifolds of the knot with Conway's notation $C(-2n,-3)$ is the same as that of the knot with Conway's notation $C(2n,3)$. For the volume formula, since the knot T_{2n} has to be hyperbolic, we exclude the case when $n = 0$.

Theorem 1.1. *Let $X_{2n}(\alpha)$, $0 \le \alpha < \alpha_0$ be the hyperbolic cone-manifold with underlying space S^3 and with singular set T_{2n} of cone-angle α. Then the volume of $X_{2n}(\alpha)$ is given by the following formula*

$$\mathrm{Vol}\left(X_{2n}(\alpha)\right) = \int_{\alpha}^{\pi} \log\left|\frac{M^{-2}+x}{M^2+x}\right| d\alpha,$$

where x with $\mathrm{Im}(x) \le 0$ is a zero of the Riley-Mednykh polynomial $P_{2n} = P_{2n}(x,M)$ which is given recursively by

$$P_{2n} = \begin{cases} QP_{2(n-1)} - M^8 P_{2(n-2)} & \text{if } n > 1, \\ QP_{2(n+1)} - M^8 P_{2(n+2)} & \text{if } n < -1, \end{cases}$$

with initial conditions

$$P_{-2} = M^2x^2 + \left(M^4 - M^2 + 1\right)x + M^2,$$
$$P_0 = M^{-2} \text{ for } n < 0 \qquad \text{and} \qquad P_0(x,M) = 1 \text{ for } n > 0,$$
$$P_2 = -M^4x^3 + \left(-2M^6 + M^4 - 2M^2\right)x^2 + \left(-M^8 + M^6 - 2M^4 + M^2 - 1\right)x + M^4,$$

and $M = e^{\frac{i\alpha}{2}}$ *and*

$$Q = -M^4x^3+\left(-2M^6+2M^4-2M^2\right)x^2+\left(-M^8+2M^6-3M^4+2M^2-1\right)x+2M^4.$$

From Theorem 1.1, the following corollary can be obtained. The following corollary gives the hyperbolic volume of the k-fold cyclic covering over T_{2n}, $M_k(X_{2n})$, for $k \geq 3$.

Corollary 1.2. *The volume of* $M_k(X_{2n})$ *is given by the following formula*

$$\mathrm{Vol}\left(M_k(X_{2n})\right)) = k\int_{\frac{2\pi}{k}}^{\pi} \log\left|\frac{M^{-2}+x}{M^2+x}\right| d\alpha,$$

where x *with* $\mathrm{Im}(x) \leq 0$ *is a zero of the Riley-Mednykh polynomial* $P_{2n} = P_{2n}(x, M)$ *and* $M = e^{i\frac{\alpha}{2}}$.

2. Two bridge knots with Conway's notation $C(2n, 3)$



Fig. 1. A two bridge knot with Conway's notation $C(2n, 3)$ for $n > 0$ (left) and for $n < 0$ (right)

A knot is a two bridge knot with Conway's notation $C(2n, 3)$ if it has a regular two-dimensional projection of the form in Figure 1. For example, Figure 2 is knot $C(4, 3)$. It has 3 left-handed vertical crossings and $2n$ right-handed horizontal crossings. Recall that we denote it by T_{2n}. In [20, Proposition 1], the fundamental group of two-bridge knots is presented. We will use the fundamental group of X_{2n} in [10]. In [10], the fundamental group of X_{2n} is calculated with 3 right-handed vertical crossings as positive crossings instead of three left-handed vertical crossings. The following proposition is tailored to our purpose. The reduced word relation of the one in the following proposition can also be obtained by reading off the fundamental group from the Schubert normal form of T_{2n} with slope $\frac{4n+1}{6n+1}$ for $n > 0$ and by taking the inverse of both sides after reading off the fundamental group from the Schubert normal form of T_{2n} with slope $\frac{4n+1}{6n+1}$ for $n < 0$ [22,20], which answers



Fig. 2. The knot $C(4,3)$ (7_3 in the Rolfsen's knot table)

the Hoste-Shanahan's question whether their presentation is actually derived from Schubert's canonical 2-bridge diagram or not.

Proposition 2.1.

$$\pi_1(X_{2n}) = \left\langle s, t \mid swt^{-1}w^{-1} = 1 \right\rangle,$$

where $w = (ts^{-1}tst^{-1}s)^n$.

3. $\left(PSL(2,C), \mathbb{H}^3\right)$ structure of $X_{2n}(\alpha)$

Let $R = \mathrm{Hom}(\pi_1(X_{2n}), \mathrm{SL}(2,\mathbb{C}))$. Given a set of generators, s, t, of the fundamental group for $\pi_1(X_{2n})$, we define a set $R(\pi_1(X_{2n})) \subset \mathrm{SL}(2,\mathbb{C})^2 \subset \mathbb{C}^8$ to be the set of all points $(\rho(s), \rho(t))$, where ρ is a representaion of $\pi_1(X_{2n})$ into $\mathrm{SL}(2,\mathbb{C})$. Since the defining relation of $\pi_1(X_{2n})$ gives the defining equation of $R(\pi_1(X_{2n}))$ [21], $R(\pi_1(X_{2n}))$ is an affine algebraic set in $\mathbb{C}^8$. $R(\pi_1(X_{2n}))$ is well-defined up to isomorphisms which arise from changing the set of generators. We say elements in R which differ by conjugations in $\mathrm{SL}(2,\mathbb{C})$ are *equivalent*. A point on the variety gives the $\left(PSL(2,C), \mathbb{H}^3\right)$ structure of $X_{2n}(\alpha)$.

Let

$$\rho(s) = \begin{bmatrix} M & 1 \\ 0 & M^{-1} \end{bmatrix}, \quad \rho(t) = \begin{bmatrix} M & 0 \\ 2 - M^2 - M^{-2} - x & M^{-1} \end{bmatrix}.$$

Then ρ becomes a representation if and only if M and x satisfies a polynomial equation [21,14]. We call the defining polynomial of the algebraic set $\{(M, x)\}$ as the *Riley-Mednykh polynomial*.

3.1. *The Riley-Mednykh polynomial*

Since we are interested in the excellent component (the geometric component) of $R(\pi_1(X_{2n}))$, in this subsection we set $M = e^{\frac{i\alpha}{2}}$. Given the fundamental group of

X_{2n},

$$\pi_1(X_{2n}) = \langle s,t \mid swt^{-1}w^{-1} = 1 \rangle,$$

where $w = (ts^{-1}tst^{-1}s)^n$, let $S = \rho(s),\ T = \rho(t)$ and $W = \rho(w)$. Then the trace of S and the trace of T are both $2\cos\frac{\alpha}{2}$.

Lemma 3.1. *For* $c \in \mathrm{SL}(2,\mathbb{C})$ *which satisfies* $cS = T^{-1}c$ *and* $c^2 = -I$,

$$SWT^{-1}W^{-1} = -(SWc)^2$$

Proof.

$$\begin{aligned}(SWc)^2 &= SWcSWc = SWT^{-1}c(TS^{-1}TST^{-1}S)^n c \\ &= SWT^{-1}(S^{-1}TS^{-1}T^{-1}ST^{-1})^n c^2 = -SWT^{-1}W^{-1}.\end{aligned}$$ □

From the structure of the algebraic set of $R(\pi_1(X_{2n}))$ with coordinates $\rho(s)$ and $\rho(t)$ we have the defining equation of $R(\pi_1(X_{2n}))$. By plugging in $e^{\frac{i\alpha}{2}}$ into M of that equation, we have the following theorem.

Theorem 3.2. *ρ is a representation of $\pi_1(X_{2n})$ if and only if x is a root of the following Riley-Mednykh polynomial $P_{2n} = P_{2n}(x,M)$ which is given recursively by*

$$P_{2n} = \begin{cases} QP_{2(n-1)} - M^8 P_{2(n-2)} & \text{if } n > 1, \\ QP_{2(n+1)} - M^8 P_{2(n+2)} & \text{if } n < -1, \end{cases}$$

with initial conditions

$P_{-2} = M^2x^2 + \left(M^4 - M^2 + 1\right)x + M^2$,

$P_0 = M^{-2}$ for $n < 0$ and $P_0(x,M) = 1$ for $n > 0$,

$P_2 = -M^4x^3 + \left(-2M^6 + M^4 - 2M^2\right)x^2 + \left(-M^8 + M^6 - 2M^4 + M^2 - 1\right)x + M^4$,

and

$Q = -M^4x^3 + \left(-2M^6 + 2M^4 - 2M^2\right)x^2 + \left(-M^8 + 2M^6 - 3M^4 + 2M^2 - 1\right)x + 2M^4$.

Proof. Note that $SWT^{-1}W^{-1} = I$, which gives the defining equations of $R(\pi_1(X_{2n}))$, is equivalent to $(SWc)^2 = -I$ in $\mathrm{SL}(2,\mathbb{C})$ by Lemma 3.1 and $(SWc)^2 = -I$ in $\mathrm{SL}(2,\mathbb{C})$ is equivalent to $\mathrm{tr}(SWc) = 0$.

We can find two c's in $\mathrm{SL}(2,\mathbb{C})$ which satisfies $cS = T^{-1}c$ and $c^2 = -I$ by direct computations. The existence and the uniqueness of the isometry (the involution) which is represented by c are shown in [4, p. 46]. Since two c's give the same element in $\mathrm{PSL}(2,\mathbb{C})$, we use one of them. Hence, we may assume

$$c = \begin{bmatrix} 0 & -\frac{M}{\sqrt{-1+2M^2-M^4-M^2x}} \\ \frac{\sqrt{-1+2M^2-M^4-M^2x}}{M} & 0 \end{bmatrix},$$

$$S = \begin{bmatrix} e^{\frac{i\alpha}{2}} & 1 \\ 0 & e^{-\frac{i\alpha}{2}} \end{bmatrix}, \quad T = \begin{bmatrix} e^{\frac{i\alpha}{2}} & 0 \\ 2 - e^{i\alpha} - e^{-i\alpha} - x\, e^{-\frac{i\alpha}{2}} & \end{bmatrix},$$

and let $U = TS^{-1}TST^{-1}S$.

Recall that P_{2n} is the defining polynomial of the algebraic set $\{(M, x)\}$ and the defining polynomial of $R(\pi_1(X_{2n}))$ corresponding to our choice of $\rho(s)$ and $\rho(t)$. We will show that P_{2n} is equal to $\mathrm{tr}(SWc)/\mathrm{tr}(Sc)$ times M^{4n} for $n > 0$ and M^{4n-2} for $n < 0$. One can easily see $\mathrm{tr}(SUc) = P_2\mathrm{tr}(Sc)/M^4$, $\mathrm{tr}(SWc) = P_{-2}\mathrm{tr}(Sc)/M^2$ and $\mathrm{tr}(U) = Q/M^4 = \mathrm{tr}(U^{-1})$. We set P_0 as the statement of the theorem. Now, we only need the following recurrence relations.

$$\begin{aligned}\mathrm{tr}(SWc) &= \mathrm{tr}(SU^nc) = \mathrm{tr}(SU^{n-1}cU^{-1}) = \mathrm{tr}(SU^{n-1}c)\mathrm{tr}(U^{-1}) - \mathrm{tr}(SU^{n-1}cU) \\ &= \mathrm{tr}(SU^{n-1}c)\mathrm{tr}(U^{-1}) - \mathrm{tr}(SU^{n-2}c) = 0 \text{ if } n > 1\end{aligned}$$

or

$$\begin{aligned}\mathrm{tr}(SWc) &= \mathrm{tr}(SU^nc) = \mathrm{tr}(SU^{n+1}cU) = \mathrm{tr}(SU^{n+1}c)\mathrm{tr}(U) - \mathrm{tr}(SU^{n+1}cU^{-1}) \\ &= \mathrm{tr}(SU^{n+1}c)\mathrm{tr}(U) - \mathrm{tr}(SU^{n+2}c) = 0 \text{ if } n < -1,\end{aligned}$$

where the third equality comes from the Cayley-Hamilton theorem. Since $\mathrm{tr}(Sc)$, $\mathrm{tr}(SUc)$, and $\mathrm{tr}(SU^{-1}c)$ have $\mathrm{tr}(Sc) = \frac{\sqrt{-1+2M^2-M^4-M^2x}}{M}$ as a common factor, all of $\mathrm{tr}(SWc)$'s have $\mathrm{tr}(Sc)$ as a common factor. We left the common factor out of $\mathrm{tr}(SWc)$, multiplied it by a power of M^{4n} if $n > 0$ and M^{4n-2} for $n < 0$ to clear the fractions and denote it by P_{2n}, the Riley-Mednykh polynomial. □

4. Longitude

Let $l = ww^*s^{-4n}$, where w^* is the word obtained by reversing w. Let $L = \rho(l)_{11}$. Then l is the longitude which is null-homologus in X_{2n}. Recall $\rho(w) = U^n$. We can write $\rho(w^*) = \widetilde{U}^n$. It is easy to see that U and $\widetilde{U}$ can be written as

$$U = \begin{pmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{pmatrix}$$

and

$$\widetilde{U} = \begin{pmatrix} \tilde{u}_{22} & \tilde{u}_{12} \\ \tilde{u}_{21} & \tilde{u}_{11} \end{pmatrix}$$

where $\tilde{u}_{ij}$ is obtained by u_{ij} by replacing M with M^{-1}. Similar computation was introduced in [10].

Definition 4.1. The *complex length* of the longitude l is the complex number γ_α modulo $4\pi\mathbb{Z}$ satisfying

$$\mathrm{tr}(\rho(l)) = 2\cosh\frac{\gamma_\alpha}{2}.$$

Note that $l_\alpha = |Re(\gamma_\alpha)|$ is the real length of the longitude of the cone-manifold $X_{2n}(\alpha)$.

The following lemma was introduced in [10] with slightly different coordinates.

Lemma 4.2. *[10]* $u_{21}L + \tilde{u}_{21}M^{-4n} = 0$.

Theorem 4.3.

$$L = -M^{-4n-2}\frac{M^{-2}+x}{M^2+x}.$$

Proof. By directly computing $u_{21}L + \tilde{u}_{21}M^{-4n} = 0$ in Lemma 4.2, the theorem follows. □

5. Proof of Theorem 1.1

For $n \geq 1$ and $M = e^{i\frac{\alpha}{2}}$, $P_{2n}(x, M)$ have $3n$ component zeros, and for $n < -1$, $-(3n+1)$ component zeros. For each n, there exists an angle $\alpha_0 \in [\frac{2\pi}{3}, \pi)$ such that $T_{2n}(\alpha)$ is hyperbolic for $\alpha \in (0, \alpha_0)$, Euclidean for $\alpha = \alpha_0$, and spherical for $\alpha \in (\alpha_0, \pi]$ [18,8,11,19]. From the following Equality (5.1), when $|L| = 1$, which happens when $\alpha = \alpha_0$, $\mathrm{Im}(x) = 0$. Hence, when α increases from 0 to α_0, two complex numbers x and $\overline{x}$ approach a same real number. In other words, $P_{2n}(x, e^{\frac{i\alpha_0}{2}})$ has a multiple root. Denote by $D(X_{2n}(\alpha))$ the discriminant of $P_{2n}(x, M)$ over x. Then α_0 will be one of the zeros of $D(X_{2n}(\alpha))$.

From Theorem 4.3, we have the following equality,

$$|L|^2 = \left|\frac{M^{-2}+x}{M^2+x}\right| = \frac{(\cos\alpha + \mathrm{Re}(x))^2 + (\mathrm{Im}(x) - \sin\alpha)^2}{(\cos\alpha + \mathrm{Re}(x))^2 + (\mathrm{Im}(x) + \sin\alpha)^2}. \tag{5.1}$$

For the volume, we choose L with $|L| \geq 1$ and hence we have $\mathrm{Im}(x) \leq 0$ by Equality (5.1). Using the Schläfli formula, we calculate the volume of $\rho(X_{2n}) = \rho(X_{2n}(0))$ for each component with $|L| \geq 1$ and having one of the zeros α_0 of $D(X_{2n}(\alpha))$ with $\alpha_0 \in [\frac{2\pi}{3}, \pi)$ on it. The component which gives the maximal volume is the excellent component [3,5]. On the geometric component we have the volume of a hyperbolic cone-manifold $X_{2n}(\alpha)$ for $0 \leq \alpha < \alpha_0$:

$$\begin{aligned}
\mathrm{Vol}(X_{2n}(\alpha)) &= -\int_{\alpha_0}^{\alpha} \frac{l_\alpha}{2}\, d\alpha \\
&= -\int_{\alpha_0}^{\alpha} \log|L|\, d\alpha \\
&= -\int_{\pi}^{\alpha} \log|L|\, d\alpha \\
&= \int_{\alpha}^{\pi} \log|L|\, d\alpha \\
&= \int_{\alpha}^{\pi} \log\left|\frac{M^{-2}+x}{M^2+x}\right|\, d\alpha,
\end{aligned}$$

where the first equality comes from the Schläfli formula for cone-manifolds (Theorem 3.20 of [1]), the second equality comes from the fact that $l_\alpha = |Re(\gamma_\alpha)|$ is the real length of the longitude of $X_{2n}(\alpha)$, the third equality comes from the fact that $\log|L| = 0$ for $\alpha_0 < \alpha \leq \pi$ by Equality 5.1 since all the characters are real (the proof of Proposition 6.4 of [19]) for $\alpha_0 < \alpha \leq \pi$, and $\alpha_0 \in [\frac{2\pi}{3}, \pi)$ is a zero of the discriminant $D(X_{2n}(\alpha))$.

Acknowledgments

The authors would like to thank Alexander Mednykh and Hyuk Kim for their various helps and anonymous referees.

References

[1] Daryl Cooper, Craig D. Hodgson, and Steven P. Kerckhoff. *Three-dimensional orbifolds and cone-manifolds*, volume 5 of *MSJ Memoirs*. Mathematical Society of Japan, Tokyo, 2000. With a postface by Sadayoshi Kojima.

[2] D. Derevnin, A. Mednykh, and M. Mulazzani. Volumes for twist link cone-manifolds. *Bol. Soc. Mat. Mexicana (3)*, 10(Special Issue):129–145, 2004.

[3] Nathan M. Dunfield. Cyclic surgery, degrees of maps of character curves, and volume rigidity for hyperbolic manifolds. *Invent. Math.*, 136(3):623–657, 1999.

[4] Werner Fenchel. *Elementary geometry in hyperbolic space*, volume 11 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1989. With an editorial by Heinz Bauer.

[5] Stefano Francaviglia and Ben Klaff. Maximal volume representations are Fuchsian. *Geom. Dedicata*, 117:111–124, 2006.

[6] Ji-Young Ham and Joongul Lee. Explicit formulae for Chern-Simons invariants of the twist knot orbifolds and edge polynomials of twist knots. `http://arxiv.org/abs/1411.2383`, 2014. Preprint.

[7] Ji-Young Ham, Alexander Mednykh, and Vladimir Petrov. Trigonometric identities and volumes of the hyperbolic twist knot cone-manifolds. *J. Knot Theory Ramifications*, 23(12):1450064, 16, 2014.

[8] Hugh Hilden, María Teresa Lozano, and José María Montesinos-Amilibia. On a remarkable polyhedron geometrizing the figure eight knot cone manifolds. *J. Math. Sci. Univ. Tokyo*, 2(3):501–561, 1995.

[9] Hugh M. Hilden, María Teresa Lozano, and José María Montesinos-Amilibia. Volumes and Chern-Simons invariants of cyclic coverings over rational knots. In *Topology and Teichmüller spaces (Katinkulta, 1995)*, pages 31–55. World Sci. Publ., River Edge, NJ, 1996.

[10] Jim Hoste and Patrick D. Shanahan. A formula for the A-polynomial of twist knots. *J. Knot Theory Ramifications*, 13(2):193–209, 2004.

[11] Sadayoshi Kojima. Deformations of hyperbolic 3-cone-manifolds. *J. Differential Geom.*, 49(3):469–516, 1998.

[12] Sadayoshi Kojima. Hyperbolic 3-manifolds singular along knots. *Chaos Solitons Fractals*, 9(4-5):765–777, 1998. Knot theory and its applications.

[13] Alexander Mednykh. The volumes of cone-manifolds and polyhedra. `http://mathlab.snu.ac.kr/~top/workshop01.pdf`, 2007. Lecture Notes, Seoul National University.

[14] Alexander Mednykh and Aleksey Rasskazov. On the structure of the canonical fundamental set for the 2-bridge link orbifolds. `www.mathematik.uni-bielefeld.de/sfb343/preprints/pr98062.ps.gz`, 1998. Universitt Bielefeld, Sonderforschungsbereich 343, Discrete Structuren in der Mathematik, Preprint, 98062.

[15] Alexander Mednykh and Alexey Rasskazov. Volumes and degeneration of cone-structures on the figure-eight knot. *Tokyo J. Math.*, 29(2):445–464, 2006.

[16] Alexander Mednykh and Andrei Vesnin. On the volume of hyperbolic Whitehead link cone-manifolds. *Sci. Ser. A Math. Sci. (N.S.)*, 8:1–11, 2002. Geometry and analysis.

[17] Alexander D. Mednykh. Trigonometric identities and geometrical inequalities for links and knots. In *Proceedings of the Third Asian Mathematical Conference, 2000 (Diliman)*, pages 352–368. World Sci. Publ., River Edge, NJ, 2002.

[18] Joan Porti. Spherical cone structures on 2-bridge knots and links. *Kobe J. Math.*, 21(1-2):61–70, 2004.

[19] Joan Porti and Hartmut Weiss. Deforming Euclidean cone 3-manifolds. *Geom. Topol.*, 11:1507–1538, 2007.

[20] Robert Riley. Parabolic representations of knot groups. I. *Proc. London Math. Soc. (3)*, 24:217–242, 1972.

[21] Robert Riley. Nonabelian representations of 2-bridge knot groups. *Quart. J. Math. Oxford Ser. (2)*, 35(138):191–208, 1984.

[22] Horst Schubert. Knoten mit zwei Brücken. *Math. Z.*, 65:133–170, 1956.